




# Visualization of Graphs

## Lecture 4:

## Straight-Line Drawings of Planar Graphs II: Schnyder Woods



Johannes Zink




# Planar Straight-Line Drawings

**Theorem.** [De Fraysseix, Pach, Pollack '90]  
 Every  $n$ -vertex planar graph has a planar straight-line drawing of size  $(2n - 4) \times (n - 2)$ .

**Theorem.** [Schnyder '90]  
 Every  $n$ -vertex planar graph has a planar straight-line drawing of size  ~~$(n - 2) \times (n - 2)$~~   $(2n - 5) \times (2n - 5)$ .

**Idea.** (easier to show)

- Fix outer triangle.
- Compute coordinates of inner vertices
  - based on outer triangle and
  - how much space there should be for other vertices
  - using weighted barycentric coordinates.





# Barycentric Coordinates

Recall:  $\text{barycenter}(x_1, \dots, x_k) = \sum_{i=1}^k x_i / k$

Let  $A, B, C$  form a triangle, and let  $x$  lie in  $\triangle ABC$ . The **barycentric coordinates** of  $x$  with respect to  $\triangle ABC$  are a triple  $(\alpha, \beta, \gamma) \in \mathbb{R}_{\geq 0}^3$  such that

- $\alpha + \beta + \gamma = 1$  and
- $x = \alpha A + \beta B + \gamma C$ .



# Barycentric Representation


A **barycentric representation** of a graph  $G = (V, E)$  is an assignment of barycentric coordinates to the vertices of  $G$ :

$$f: V \rightarrow \mathbb{R}_{\geq 0}^3, v \mapsto (v_1, v_2, v_3)$$

with the following properties:

(B1)  $v_1 + v_2 + v_3 = 1$  for all  $v \in V$ ,

(B2) for each  $\{x, y\} \in E$  and each  $z \in V \setminus \{x, y\}$  there exists a  $k \in \{1, 2, 3\}$  with  $x_k < z_k$  and  $y_k < z_k$ .



# Barycentric Representations of Planar Graphs

## Lemma.

Let  $f: v \mapsto (v_1, v_2, v_3)$  be a barycentric representation of a planar graph  $G$ , and let  $A, B, C \in \mathbb{R}^2$  be in general position. Then the mapping

$$\phi: v \in V \mapsto v_1 A + v_2 B + v_3 C$$

yields a **planar straight-line** drawing of  $G$  inside  $\triangle ABC$ .


- No vertex  $x$  can lie on an edge  $\{u, v\}$ . (clear by definition)
- No pair of edges  $\{u, v\}$  and  $\{u', v'\}$  crosses:

$$u'_i > u_i, v_i \quad v'_j > u_j, v_j \quad u_k > u'_k, v'_k \quad v_l > u'_l, v'_l$$

$$\Rightarrow \{i, j\} \cap \{k, l\} = \emptyset$$

w.l.o.g.  $i = j = 2 \Rightarrow u'_2, v'_2 > u_2, v_2 \Rightarrow$  separated by a straight line


no three points  
on a line



How to find a  
barycentric  
representation?

# Schnyder Labeling


Let  $\phi: v \mapsto (v_1, v_2, v_3)$  be a barycentric representation of a planar graph  $G$ , and let  $A, B, C \in \mathbb{R}^2$  be in general position.



# Schnyder Labeling

Let  $\phi: v \mapsto (v_1, v_2, v_3)$  be a barycentric representation of a planar graph  $G$ , and let  $A, B, C \in \mathbb{R}^2$  be in general position.

We can label each angle in  $\triangle xyz$  **uniquely** with  $k \in \{1, 2, 3\}$ .



# Schnyder Labeling

Let  $\phi: v \mapsto (v_1, v_2, v_3)$  be a barycentric representation of a planar graph  $G$ , and let  $A, B, C \in \mathbb{R}^2$  be in general position.



We can label each angle in  $\triangle xyz$  **uniquely** with  $k \in \{1, 2, 3\}$ .

A **Schnyder labeling** of a plane triangulation  $G$  is a labeling of all internal angles with labels **1**, **2** and **3** such that:

**Faces:** The three angles of an internal face are labeled **1**, **2** and **3** in counterclockwise (ccw) order.

**Vertices:** The ccw order of labels around each vertex consists of

- a non-empty interval of **1**s
- followed by a non-empty interval of **2**s
- followed by a non-empty interval of **3**s.






# Schnyder Wood



A Schnyder labeling induces an edge labeling.

A **Schnyder wood** (or **realizer**) of a plane triangulation  $G = (V, E)$  is a partition of the inner edges of  $E$  into three sets of oriented edges  $T_1$ ,  $T_2$ ,  $T_3$  such that, for each inner vertex  $v \in V$ , it holds that


- $v$  has one outgoing edge in each of  $T_1$ ,  $T_2$ , and  $T_3$ .
- The ccw order of edges around  $v$  is:  
 leaving in  $T_1$ , entering in  $T_3$ , leaving in  $T_2$ ,  
 entering in  $T_1$ , leaving in  $T_3$ , entering in  $T_2$ .



# Schnyder Wood – Example and Properties





# Schnyder Wood – Example and Properties



(a Schnyder labeling is not unique)

# Schnyder Wood – Example and Properties




- All inner edges incident to  $a$ ,  $b$ , and  $c$  are incoming in the same color.
- $T_1$ ,  $T_2$ , and  $T_3$  are trees. Each spans all inner vertices and one outer vertex (its root).

# Schnyder Wood – Existence

**Lemma.**

[Kampen 1976]

Let  $G$  be a plane triangulation with vertices  $a, b, c$  on the outer face. Then there exists a **contractible edge**  $\{a, x\}$  in  $G$  with  $x \notin \{b, c\}$ .



... requires that  $a$  and  $x$  have exactly two common neighbors.

# Schnyder Wood – Existence

## Lemma.


[Kampen 1976]

Let  $G$  be a plane triangulation with vertices  $a, b, c$  on the outer face. Then there exists a **contractible edge**  $\{a, x\}$  in  $G$  with  $x \notin \{b, c\}$ .

## Theorem.

Every plane triangulation has a Schnyder labeling and a Schnyder wood.

**Proof** by induction on  $\#$  vertices via edge contractions.




This constructive proof yields an algorithm for computing a Schnyder labeling. It can be implemented to run in  $\mathcal{O}(n)$  time.

... requires that  $a$  and  $x$  have exactly two common neighbors.

→ *Exercise* 😊

# Schnyder Wood – More Properties




- From each vertex  $v$  there exists a directed red path  $P_1(v)$  to  $a$ , a directed blue path  $P_2(v)$  to  $b$ , and a directed green path  $P_3(v)$  to  $c$ .

$P_i(v)$ : path from  $v$  to root of  $T_i$ .

## Lemma.

- $P_1(v)$ ,  $P_2(v)$ ,  $P_3(v)$  cross only at  $v$ .

# Schnyder Wood – More Properties



- From each vertex  $v$  there exists a directed **red** path  $P_1(v)$  to  $a$ , a directed **blue** path  $P_2(v)$  to  $b$ , and a directed **green** path  $P_3(v)$  to  $c$ .

$P_i(v)$ : path from  $v$  to root of  $T_i$ .

$R_1(v)$ : set of faces contained in  $P_2, bc, P_3$ .

$R_2(v)$ : set of faces contained in  $P_3, ca, P_1$ .


$R_3(v)$ : set of faces contained in  $P_1, ab, P_2$ .

## Lemma.

- $P_1(v), P_2(v), P_3(v)$  cross only at  $v$ .
- For inner vertices  $u \neq v$  it holds that  $u \in R_i(v) \Rightarrow R_i(u) \subsetneq R_i(v)$ .



# Schnyder Wood – More Properties



- From each vertex  $v$  there exists a directed red path  $P_1(v)$  to  $a$ , a directed blue path  $P_2(v)$  to  $b$ , and a directed green path  $P_3(v)$  to  $c$ .

$P_i(v)$ : path from  $v$  to root of  $T_i$ .

$R_1(v)$ : set of faces contained in  $P_2, bc, P_3$ .


$R_2(v)$ : set of faces contained in  $P_3, ca, P_1$ .

$R_3(v)$ : set of faces contained in  $P_1, ab, P_2$ .

## Lemma.

- $P_1(v), P_2(v), P_3(v)$  cross only at  $v$ .
- For inner vertices  $u \neq v$  it holds that  $u \in R_i(v) \Rightarrow R_i(u) \subsetneq R_i(v)$ .

# Schnyder Wood – More Properties



- From each vertex  $v$  there exists a directed **red** path  $P_1(v)$  to  $a$ , a directed **blue** path  $P_2(v)$  to  $b$ , and a directed **green** path  $P_3(v)$  to  $c$ .

$P_i(v)$ : path from  $v$  to root of  $T_i$ .

$R_1(v)$ : set of faces contained in  $P_2, bc, P_3$ .

$R_2(v)$ : set of faces contained in  $P_3, ca, P_1$ .

$R_3(v)$ : set of faces contained in  $P_1, ab, P_2$ .

## Lemma.

- $P_1(v), P_2(v), P_3(v)$  cross only at  $v$ .
- For inner vertices  $u \neq v$  it holds that  $u \in R_i(v) \Rightarrow R_i(u) \subsetneq R_i(v)$ .
- $|R_1(v)| + |R_2(v)| + |R_3(v)| = 2n - 5$

# Schnyder Drawing

## Theorem.


[Schnyder '90]

For a plane triangulation  $G$ , the mapping

$$f: v \mapsto (v_1, v_2, v_3) = \frac{1}{2n-5} (|R_1(v)|, |R_2(v)|, |R_3(v)|)$$

is a barycentric representation of  $G$  and, thus, yields a planar straight-line drawing of  $G$

- (B1)  $v_1 + v_2 + v_3 = 1$  for all  $v \in V$  ✓
- (B2) for each  $\{x, y\} \in E$  and each  $z \in V \setminus \{x, y\}$   
there exists  $k \in \{1, 2, 3\}$  with  $x_k < z_k$  and  $y_k < z_k$ 
  - $\{x, y\}$  must lie in  $R_i(z)$  for some  $i \in \{1, 2, 3\}$



# Schnyder Drawing

Set  $A = (0, 0)$ ,  $B = (2n - 5, 0)$ , and  $C = (0, 2n - 5)$ .

## Theorem.


[Schnyder '90]

For a plane triangulation  $G$ , the mapping



$$f: v \mapsto (v_1, v_2, v_3) = \frac{1}{2n-5} (|R_1(v)|, |R_2(v)|, |R_3(v)|)$$

is a barycentric representation of  $G$  and, thus, yields a planar straight-line drawing of  $G$  on the  $(2n - 5) \times (2n - 5)$  grid.

- (B1)  $v_1 + v_2 + v_3 = 1$  for all  $v \in V$  ✓
- (B2) for each  $\{x, y\} \in E$  and each  $z \in V \setminus \{x, y\}$   
there exists  $k \in \{1, 2, 3\}$  with  $x_k < z_k$  and  $y_k < z_k$  ✓
- $\{x, y\}$  must lie in  $R_i(z)$  for some  $i \in \{1, 2, 3\}$
  - $x, y \in R_i(z) \Rightarrow R_i(x), R_i(y) \subsetneq R_i(z)$   
 $\Rightarrow |R_i(x)|, |R_i(y)| < |R_i(z)|$





# Schnyder Drawing – Example



- $n = 7; \quad 2n - 5 = 9$
- $f(v_1) = (9, 0, 0)$
- $f(v_2) = (0, 9, 0)$
- $f(v_3) = (2, 6, 1)$
- $f(v_4) = (5, 2, 2)$
- $f(v_5) = (1, 2, 6)$
- $f(v_6) = (4, 1, 4)$
- $f(v_7) = (0, 0, 9)$

# Schnyder Drawing – Example



$n = 7;$	$2n - 5 = 9$	$f(v_4) = (5, 2, 2)$
$f(v_1) = (9, 0, 0)$		$f(v_5) = (1, 2, 6)$
$f(v_2) = (0, 9, 0)$		$f(v_6) = (4, 1, 4)$
$f(v_3) = (2, 6, 1)$		$f(v_7) = (0, 0, 9)$

# Weak Barycentric Representation

A **weak barycentric representation** of a graph  $G = (V, E)$  is an assignment of barycentric coordinates to  $V$ :

$$\phi: V \rightarrow \mathbb{R}_{\geq 0}^3, v \mapsto (v_1, v_2, v_3)$$

with the following properties:

(W1)  $v_1 + v_2 + v_3 = 1$  for all  $v \in V$ ,

(W2) for each  $\{x, y\} \in E$  and each  $z \in V \setminus \{x, y\}$  there exists a  $k \in \{1, 2, 3\}$  with


$$(x_k, x_{k+1}) <_{\text{lex}} (z_k, z_{k+1}) \text{ and } (y_k, y_{k+1}) <_{\text{lex}} (z_k, z_{k+1}).$$

## Lemma.

For a weak barycentric representation  $\phi: v \mapsto (v_1, v_2, v_3)$  and a triangle  $\triangle ABC$ , the mapping

$$f: v \in V \mapsto v_1A + v_2B + v_3C$$

yields a **planar** drawing of  $G$  inside  $\triangle ABC$ .




i.e., either  $y_k < z_k$  or  
 $y_k = z_k$  and  $y_{k+1} < z_{k+1}$

indices modulo 3

**Proof.**  $\rightarrow$  *Exercise!*

# Counting Vertices



$P_i(v)$ : path from  $v$  to root of  $T_i$ .

$R_1(v)$ : subgraph bounded by  $P_2$ ,  $bc$ ,  $P_3$ .


$R_2(v)$ : subgraph bounded by  $P_3$ ,  $ca$ ,  $P_1$ .

$R_3(v)$ : subgraph bounded by  $P_1$ ,  $ab$ ,  $P_2$ .

$$v_i = |V(R_i(v))| - |P_{i-1}(v)|$$



# Counting Vertices



$P_i(v)$ : path from  $v$  to root of  $T_i$ .

$R_1(v)$ : subgraph bounded by  $P_2$ ,  $bc$ ,  $P_3$ .

$R_2(v)$ : subgraph bounded by  $P_3$ ,  $ca$ ,  $P_1$ .

$R_3(v)$ : subgraph bounded by  $P_1$ ,  $ab$ ,  $P_2$ .


$$v_i = |V(R_i(v))| - |P_{i-1}(v)|$$

$$v_1 = 10 - 3 = 7$$

$$v_2 = 6 - 3 = 3$$

$$v_3 = 8 - 3 = 5$$

# Counting Vertices



$P_i(v)$ : path from  $v$  to root of  $T_i$ .

$R_1(v)$ : subgraph bounded by  $P_2$ ,  $bc$ ,  $P_3$ .

$R_2(v)$ : subgraph bounded by  $P_3$ ,  $ca$ ,  $P_1$ .

$R_3(v)$ : subgraph bounded by  $P_1$ ,  $ab$ ,  $P_2$ .

$$v_i = |V(R_i(v))| - |P_{i-1}(v)|$$

$$v_1 = 10 - 3 = 7$$


$$v_2 = 6 - 3 = 3$$

$$v_3 = 8 - 3 = 5$$

## Lemma.

- For inner vertices  $u \neq v$  it holds that  $u \in R_i(v) \Rightarrow (u_i, u_{i+1}) <_{\text{lex}} (v_i, v_{i+1})$ .

# Counting Vertices



$P_i(v)$ : path from  $v$  to root of  $T_i$ .

$R_1(v)$ : subgraph bounded by  $P_2$ ,  $bc$ ,  $P_3$ .

$R_2(v)$ : subgraph bounded by  $P_3$ ,  $ca$ ,  $P_1$ .

$R_3(v)$ : subgraph bounded by  $P_1$ ,  $ab$ ,  $P_2$ .

$$v_i = |V(R_i(v))| - |P_{i-1}(v)|$$

$$v_1 = 10 - 3 = 7$$

$$v_2 = 6 - 3 = 3$$

$$v_3 = 8 - 3 = 5$$

## Lemma.

■ For inner vertices  $u \neq v$  it holds that  $u \in R_i(v) \Rightarrow (u_i, u_{i+1}) <_{\text{lex}} (v_i, v_{i+1})$ .

■  $v_1 + v_2 + v_3 = n - 1$

# Schnyder Drawing<sup>\*</sup>

Set  $A = (0, 0)$ ,  $B = (n - 1, 0)$ , and  $C = (0, n - 1)$ .

## Theorem.



[Schnyder '90]

For a plane triangulation  $G$ , the mapping

$$f: v \mapsto \frac{1}{n-1}(v_1, v_2, v_3)$$

is a barycentric representation of  $G$  and, thus, yields a planar straight-line drawing of  $G$  on the  $(n - 2) \times (n - 2)$  grid.

# Schnyder Drawing\* – Example



$$n = 16, n - 2 = 14$$

$$f(a) = (n - 2, 1, 0)$$

# Results & Variations

## Theorem.



[De Fraysseix, Pach, Pollack '90]

Every  $n$ -vertex planar graph has a planar straight-line drawing of size  $(2n - 4) \times (n - 2)$ . Such a drawing can be computed in  $O(n)$  time.

## Theorem.

[Schnyder '90]

Every  $n$ -vertex planar graph has a planar straight-line drawing of size  $(n - 2) \times (n - 2)$ . Such a drawing can be computed in  $O(n)$  time.



# Results & Variations

## Theorem.

[De Fraysseix, Pach, Pollack '90]

Every  $n$ -vertex planar graph has a planar straight-line drawing of size  $(2n - 4) \times (n - 2)$ . Such a drawing can be computed in  $O(n)$  time.

## Theorem.

[Schnyder '90]

Every  $n$ -vertex planar graph has a planar straight-line drawing of size  $(n - 2) \times (n - 2)$ . Such a drawing can be computed in  $O(n)$  time.



*Exercise!*

## Theorem.

[Brandenburg '08]

Every  $n$ -vertex planar graph has a planar straight-line drawing of size  $\frac{4}{3}n \times \frac{2}{3}n$ . Such a drawing can be computed in  $O(n)$  time.

# Results & Variations





# Results & Variations

## Theorem.

[Kant '96]

Every  $n$ -vertex 3-connected planar graph has a planar straight-line drawing of size  $(2n - 4) \times (n - 2)$  where all faces are drawn convex. Such a drawing can be computed in  $O(n)$  time.

## Theorem.

[Chrobak & Kant '97]

Every  $n$ -vertex 3-connected planar graph has a planar straight-line drawing of size  $(n - 2) \times (n - 2)$  where all faces are drawn convex. Such a drawing can be computed in  $O(n)$  time.

## Theorem.

[Felsner '01]

Every 3-connected planar graph with  $f$  faces has a planar straight-line drawing of size  $(f - 1) \times (f - 1)$  where all faces are drawn convex. Such a drawing can be computed in  $O(n)$  time.

# Literature

- [PGD Ch. 4.3] for detailed explanation of Schnyder woods etc.
- [Sch90] “Embedding planar graphs on the grid”, Walter Schnyder, SoCG 1990 – original paper on Schnyder realizer method.