



# Visualization of Graphs

## Lecture 7:

### Contact Representations of Planar Graphs: Triangle Contacts and Rectangular Duals



Johannes Zink




# Intersection Representation of Graphs

In an **intersection representation** of a graph,


- each vertex is represented by a set
- such that two sets intersect  $\Leftrightarrow$  the corresponding vertices are adjacent.

For a collection  $\mathcal{S}$  of sets, the **intersection graph**  $G(\mathcal{S})$  of  $\mathcal{S}$  has vertex set  $\mathcal{S}$  and edge set  $\{\{S, S'\} : S, S' \in \mathcal{S}, S \neq S', \text{ and } S \cap S' \neq \emptyset\}$ .







# Contact Representation of Graphs

Let  $G$  be a graph.




Let  $\mathcal{S}$  be a family of geometric objects (e.g., disks).




Represent each vertex  $v$  by a geometric object  $S(v) \in \mathcal{S}$



rectangular cuboids



In an  **$\mathcal{S}$ -contact representation** of  $G$ ,  $S(u)$  and  $S(v)$  touch iff  $uv \in E$



$G$  is planar

[Koebe 1936]

disks

polygons



A contact representation is an intersection representation with interior-disjoint sets.

# Contact Representation of Planar Graphs


Is the intersection graph of a contact representation always planar?

- No, not even for connected object types in the plane.


Some object types are used to represent **special classes** of planar graphs:



bipartite planar graphs



max. triangle-free planar graphs




planar triangulations

# General Approach

How to compute a contact representation of a given graph  $G$ ?


- Consider only inner triangulations (or maximal bipartite graphs, etc.)
  - Triangulate by adding vertices, not by adding edges
- Describe contact representation combinatorially.
  - Which objects touch each other in which way?
- Compute combinatorial description.
- Show that combinatorial description can be used to construct drawing.



# This Lecture


Representation with right-triangles and corner contact:

- Use Schnyder realizer to describe contacts between triangles.
- Use canonical order to compute drawing.



Representation with dissection of a rectangle, called **rectangular dual**:

- Find a description similar to a Schnyder realizer for rectangles.
- Construct drawing via *st*-digraphs, duals, and topological sorting.




# Triangle Corner Contact Representation

## Main Idea.


Use canonical order and Schnyder realizer to find coordinates for triangles.

## Detailed Idea.

- Place base of triangle at height equal to position in canonical order.
- Triangle tip is precisely at base of triangle corresponding to cover neighbor.
- Outgoing edges in Schnyder forest indicate corner contacts.




# Triangle Contact Representation Example







# Triangle Contact Representation Example




# Triangle Contact Representation Example




# T-shape Contact Representation




# T-shape Contact Representation





# Cartograms




© worldmapper.org






© New York Times



© Bettina Speckmann




© New York Times




# Rectangular Dual

Exactly 4 vertices on outer face





PTP

Properly Triangulated  
Planar Graph  $G$



RD


Rectangular Dual  $\mathcal{R}$



no separating  
triangle

A **rectangular dual** of a graph  $G$  is a contact representation with axis-aligned rectangles s.t.

- no four rectangles share a point, and
- the union of all rectangles is a rectangle




## Theorem.


A graph  $G$  has a rectangular dual  $\mathcal{R}$  if and only if  $G$  is a PTP graph.

[Kozłmiński, Kinnen '85]


# Regular Edge Labeling




Properly Triangulated  
Planar Graph  $G$



Rectangular Dual  $\mathcal{R}$




# Regular Edge Labeling




PTP

Properly Triangulated  
Planar Graph  $G$




RD

Rectangular Dual  $\mathcal{R}$







# Regular Edge Labeling




Properly Triangulated  
Planar Graph  $G$



Rectangular Dual  $\mathcal{R}$




# Regular Edge Labeling




Properly Triangulated  
Planar Graph  $G$

PTP





Regular Edge Labeling

REL




Rectangular Dual  $\mathcal{R}$

RD




for every  
inner vertex




for four  
outer vertices

[Kant, He '94]: In linear time



PTP

$O(n)$



REL

$O(n)$




RD

# Refined Canonical Order


## Theorem.

Let  $G$  be a PTP graph. There exists a labeling  $v_1 = v_S, v_2 = v_W, v_3, \dots, v_n = v_N$  of the vertices of  $G$  such that for every  $4 \leq k \leq n$ :


- The subgraph  $G_{k-1}$  induced by  $v_1, \dots, v_{k-1}$  is biconnected and the boundary  $C_{k-1}$  of  $G_{k-1}$  contains the edge  $(v_S, v_W)$ .
- $v_k$  is in exterior face of  $G_{k-1}$ , and its neighbors in  $G_{k-1}$  form an (at least 2-element) subinterval of the path  $C_{k-1} \setminus (v_S, v_W)$ .
- If  $k \leq n - 2$ , then  $v_k$  has at least 2 neighbors in  $G \setminus G_{k-1}$ .




# Refined Canonical Order Example




# Refined Canonical Order Example



# Refined Canonical Order Example




# Refined Canonical Order Example



# Refined Canonical Order $\rightarrow$ REL

We construct a REL as follows:

- For  $i < j$ , orient  $(v_i, v_j)$  from  $v_i$  to  $v_j$ ;
- $v_k$  has incoming edges from  $v_{b_1}, \dots, v_{b_l}$ , we say that  $v_{b_1}$  is **left point** of  $v_k$  and  $v_{b_l}$  is **right point** of  $v_k$ .
- **Base edge** of  $v_k$  is  $(v_{b_a}, v_k)$ , where  $b_a \in \{b_1, \dots, b_l\}$  is minimal.
- If  $v_{t_1}, \dots, v_{t_o}$  are higher numbered neighbors of  $v_k$ , we call  $(v_k, v_{t_1})$  **left edge** and  $(v_k, v_{t_o})$  **right edge**.



## Lemma 1.

A left edge or right edge cannot be a base edge.

**Proof.** Suppose left edge  $(v_k, v_{t_1})$  is base edge of  $v_{t_1}$ .  
 Since  $G$  triangulated,  $(v_{b_1}, v_{t_1}) \in E(G)$ .  
 Contradiction since  $k > b_1$ .




# Refined Canonical Order $\rightarrow$ REL

## Lemma 2.

An edge is either a **left edge**, a **right edge** or a **base edge**.

## Proof.

- Exclusive “or” follows from Lemma 1.
- Let  $(v_{b_a}, v_k)$  be **base edge** of  $v_k$ .
- $v_{b_a}$  is **right point** of  $v_{b_{a-1}}$ .
  - $v_{b_i}$  has at least two higher-numbered neighbors.
  - One of them is  $v_k$ ; the other one is  $v_{b_{i-1}}$  or  $v_{b_{i+1}}$ .
  - For  $1 \leq i < a - 1$ , it is  $v_{b_{i-1}}$ . Thus,  $v_{b_i}$  is right point of  $v_{b_{i-1}}$ .
- Analogously,  $v_{b_i}$  is **left point** of  $v_{b_{i+1}}$  for  $i \geq a$ .
- Edges  $(v_{b_i}, v_k)$ ,  $1 \leq i < a - 1$ , are **right edges**.
- Similarly,  $(v_{b_i}, v_k)$ , for  $a + 1 \leq i \leq l$ , are **left edges**.



# Refined Canonical Order $\rightarrow$ REL

## Coloring.

- Color **right** (**left**) edges in **red** (**blue**).
- Color a **base edge**  $(v_{b_i}, v_k)$  **red** if  $i = 1$  and **blue** if  $i = l$  and otherwise arbitrarily.



Let  $T_r$  be the red edges and  $T_b$  the blue edges.

### Lemma 3.

$\{T_r, T_b\}$  is a regular edge labeling.


## Proof.

$$t_o \geq 2$$




- $t_1 < t_2 < \dots < t_d$  and  $t_d > t_{d+1} > \dots > t_o$
  - $(v_k, v_{t_i}), 2 \leq i \leq d-1$  are **blue**
  - $(v_k, v_{t_i}), d+1 \leq i \leq o-1$  are **red**
  - $(v_k, v_{t_d})$  is either **red** or **blue**
- $\Rightarrow$  Circular order of outgoing edges at  $v_k$  correct.


# From REL to *st*-Digraphs to Coordinates




# From REL to $st$ -Digraphs to Coordinates




# From REL to *st*-Digraphs to Coordinates




SN network  $G_{ver}$   
 dual of  $G_{ver}$



# From REL to *st*-Digraphs to Coordinates



# From REL to *st*-Digraphs to Coordinates




dual of  $G_{\text{ver}}$

compute topological order


i.e., linear order of all vertices such that all directed edges point in the same direction

# From REL to *st*-Digraphs to Coordinates






# From REL to *st*-Digraphs to Coordinates



# From REL to $st$ -Digraphs to Coordinates




# Rectangular Dual Algorithm

For a PTP graph  $G = (V, E)$ :

- Find a REL  $\{T_r, T_b\}$  of  $G$ ;
- Construct a SN network  $G_{\text{ver}}$  of  $G$  (consists of  $T_b$  plus outer edges)
- Construct the dual  $G_{\text{ver}}^*$  of  $G_{\text{ver}}$  and compute a topological ordering  $f_{\text{ver}}$  of  $G_{\text{ver}}^*$
- For each vertex  $v \in V$ , let  $g$  and  $h$  be the face on the left and face on the right of  $v$ . Set  $x_1(v) = f_{\text{ver}}(g)$  and  $x_2(v) = f_{\text{ver}}(h)$ .
- Define  $x_1(v_N) = 0$ ,  $x_1(v_S) = 1$  and  $x_2(v_N) = \max f_{\text{ver}} - 1$ ,  $x_2(v_S) = \max f_{\text{ver}}$
- Analogously compute  $y_1$  and  $y_2$  with  $G_{\text{hor}}$ .
- For each  $v \in V$ , let  $R(v) = [x_1(v), x_2(v)] \times [y_1(v), y_2(v)]$ .

# Reading off Coordinates to Get Rectangular Dual



$$x_1(v_N) = 0, x_2(v_N) = 15$$

$$x_1(v_S) = 1, x_2(v_S) = 16$$

$$x_1(v_W) = 0, x_2(v_W) = 1$$

$$x_1(v_E) = 15, x_2(v_E) = 16$$

$$x_1(a) = 1, x_2(a) = 3$$

$$x_1(b) = 3, x_2(b) = 5$$

$$x_1(c) = 5, x_2(c) = 14$$

$$x_1(d) = 14, x_2(d) = 15$$

$$x_1(e) = 13, x_2(e) = 15$$

...

$$y_1(v_W) = 0, y_2(v_W) = 9$$

$$y_1(v_E) = 1, y_2(v_E) = 10$$

$$y_1(v_N) = 9, y_2(v_N) = 10$$


$$y_1(v_S) = 0, y_2(v_S) = 1$$

$$y_1(a) = 1, y_2(a) = 2$$

$$y_1(b) = 1, y_2(b) = 2$$

...

# Reading off Coordinates to Get Rectangular Dual



$$x_1(v_N) = 0, \quad x_2(v_N) = 15$$

$$x_1(v_S) = 1, \quad x_2(v_S) = 16$$

$$x_1(v_W) = 0, \quad x_2(v_W) = 1$$

$$x_1(v_E) = 15, \quad x_2(v_E) = 16$$

$$x_1(a) = 1, \quad x_2(a) = 3$$

$$x_1(b) = 3, \quad x_2(b) = 5$$

$$x_1(c) = 5, \quad x_2(c) = 14$$

$$x_1(d) = 14, \quad x_2(d) = 15$$

$$x_1(e) = 13, \quad x_2(e) = 15$$

...

$$y_1(v_W) = 0, \quad y_2(v_W) = 9$$

$$y_1(v_E) = 1, \quad y_2(v_E) = 10$$

$$y_1(v_N) = 9, \quad y_2(v_N) = 10$$

$$y_1(v_S) = 0, \quad y_2(v_S) = 1$$


$$y_1(a) = 1, \quad y_2(a) = 2$$

$$y_1(b) = 1, \quad y_2(b) = 2$$

...


# Correctness of Algorithm (Sketch)

- If edge  $(u, v)$  exists, then  $x_2(u) = x_1(v)$



$$x_2(u) = f_{\text{ver}}(g) = x_1(v)$$

- and the vertical segments of their rectangles overlap



$$\begin{aligned} y_1(v) = f_{\text{hor}}(a) &\leq y_1(u) = f_{\text{hor}}(b) \\ &< y_2(v) = f_{\text{hor}}(c) &\leq y_2(u) = f_{\text{hor}}(d) \end{aligned}$$

- If path from  $u$  to  $v$  in red at least two edges long, then  $x_2(u) < x_1(v)$ .
- No two boxes overlap.
- For details, see He's paper [He '93].

# Rectangular Dual Result

## Theorem.

Every PTP graph  $G$  has a rectangular dual, which can be computed in linear time.


## Proof.

- Compute a planar embedding of  $G$ .
- Compute a refined canonical ordering of  $G$ .
- Traverse the graph and color the edges.  $\rightarrow$  REL
- Construct  $G_{\text{ver}}$  and  $G_{\text{hor}}$ .
- Construct their duals  $G_{\text{ver}}^*$  and  $G_{\text{hor}}^*$ .
- Compute a topological ordering for vertices of  $G_{\text{ver}}^*$  and  $G_{\text{hor}}^*$ .
- Assigning coordinates to the rectangles representing vertices.


# Discussion

- A layout is **area-universal** if any assignment of areas to rectangles can be realized by a combinatorially equivalent rectangular layout.
- A rectangular layout is **area-universal** if and only if it is **one-sided**.  
[Eppstein et al. SIAM J. Comp. 2012]

one-sided




$s$



not one-sided

i.e., every segment belongs to exactly one rectangle

- Area-universal **rectilinear** representation: possible for all planar graphs.
- [Alam et al. 2013]: 8 sides (matches the lower bound)





# Literature

Construction of triangle contact representations based on

- [de Fraysseix, Ossona de Mendez, Rosenstiehl '94] On Triangle Contact Graphs

Construction of rectangular dual based on

- [He '93] On Finding the Rectangular Duals of Planar Triangulated Graphs
- [Kant, He '94] Two algorithms for finding rectangular duals of planar graphs

and originally from

- [Kozłmiński, Kinnen '85] Rectangular Duals of Planar Graphs