


Visualization of Graphs


Lecture 8:

Contact Representations of Planar Graphs: Triangle Contacts and Rectangular Duals



Part I: Geometric Representations


Jonathan Klawitter



Intersection Representation

In an **intersection representation** of a graph each vertex is represented as a set such that two sets intersect if and only if the corresponding vertices are adjacent.


For a collection \mathcal{S} of sets S_1, \dots, S_n , the **intersection graph** $G(\mathcal{S})$ of \mathcal{S} has vertex set \mathcal{S} and edge set $\{S_i S_j : i, j \in \{1, \dots, n\}, i \neq j, \text{ and } S_i \cap S_j \neq \emptyset\}$.



Contact Representation of Graphs





A contact representation is an intersection representation with interior-disjoint sets.

Let G be a graph.




Let \mathcal{S} be a set of geometric objects




Represent each vertex v by a geometric object $S(v) \in \mathcal{S}$



rectangular cuboids



In an \mathcal{S} contact representation of G , $S(u)$ and $S(v)$ touch iff $uv \in E$



G is planar

→ [Koebe 1936]

disks

→


polygons

Contact Representation of Planar Graphs


Is the intersection graph of a contact representation always planar?

- No, not even for connected object types.


Some object types are used to represent **special classes** of planar graphs:




bipartite graphs



max. triangle-free graphs




planar triangulations



General Approach

How to compute a contact representation of a given graph G ?


- Consider only inner triangulations (or maximally bipartite graphs, etc)
 - Triangulate by adding vertices, not by adding edges
- Describe contact representation combinatorically.
 - Which objects contact each other in which way?
- Compute combinatorial description.
- Show that combinatorial description can be used to construct drawing.



In This Lecture


Representations with right-triangles and corner contact

- Use Schnyder realizer to describe contacts between triangles
- Use canonical order to calculate drawing



Representation with dissection of a rectangle, called **rectangular dual**


- Find similar description like Schnyder realizer for rectangles
- Construct drawing via st-digraphs, duals, and topological sorting



Visualization of Graphs


Lecture 8:

Contact Representations of Planar Graphs: Triangle Contacts and Rectangular Duals



Part II: Triangle Contact Representations

Jonathan Klawitter




Triangle Corner Contact Representation

Idea.


Use canonical order and Schnyder realizer to find coordinates for triangles.

Observation.


- Can set base of triangle at height equal to position in canonical order.
- Triangle tip is precisely at base of triangle corresponding to cover neighbor.
- Outgoing edges in Schnyder forest indicate corner contacts.




Triangle Contact Representation Example




Triangle Contact Representation Example




Triangle Contact Representation Example



T-shape Contact Representation




T-shape Contact Representation



Visualization of Graphs


Lecture 8:

Contact Representations of Planar Graphs: Triangle Contacts and Rectangular Duals




Part III: Rectangular Duals


Jonathan Klawitter





Cartograms




© worldmapper.org






© New York Times



© Bettina Speckmann




© New York Times




Rectangular Dual

Exactly 4 vertices on outer face





PTP

Properly Triangulated
Planar Graph G



RD


Rectangular Dual \mathcal{R}



no separating
triangle

A **rectangular dual** of a graph G is a contact representation with axis aligned rectangles s.t.

- no four rectangles share a point, and
- the union of all rectangles is a rectangle




Theorem.

A graph G has a rectangular dual \mathcal{R} if and only if G is a PTP graph.


[Kozłowski, Kinnen '85]

Regular Edge Labeling




PTP

Properly Triangulated
Planar Graph G




RD

Rectangular Dual \mathcal{R}




Regular Edge Labeling




PTP

Properly Triangulated
Planar Graph G




RD

Rectangular Dual \mathcal{R}




Regular Edge Labeling




Properly Triangulated
Planar Graph G

PTP




Rectangular Dual \mathcal{R}

RD




Regular Edge Labeling




PTP

Properly Triangulated
Planar Graph G





REL

Regular Edge Labeling




RD


Rectangular Dual \mathcal{R}




Regular Edge Labeling





Properly Triangulated
Planar Graph G




Regular Edge Labeling



Rectangular Dual \mathcal{R}




for every
inner vertex




for four
outer vertices


Regular Edge Labeling




Properly Triangulated
Planar Graph G




Regular Edge Labeling




Rectangular Dual \mathcal{R}





[Kant, He '94]: In linear time




$O(n)$



$O(n)$




[Kant, He '94]:



Visualization of Graphs


Lecture 8:

Conact Representations of Planar Graphs: Triangle Contacts and Rectangular Duals



Part IV: Computing a REL

Jonathan Klawitter




Refined Canonical Order


Theorem.

Let G be a PTP graph. There exists a labeling $v_1 = v_S, v_2 = v_W, v_3, \dots, v_n = v_N$ of the vertices of G such that for every $4 \leq k \leq n$:


- The subgraph G_{k-1} induced by v_1, \dots, v_{k-1} is biconnected and boundary C_{k-1} of G_{k-1} contains the edge (v_S, v_W) .
- v_k is in exterior face of G_{k-1} , and its neighbors in G_{k-1} form a (at least 2-element) subinterval of the path $C_{k-1} \setminus (v_S, v_W)$.
- If $k \leq k - 2$, then v_k has at least 2 neighbors in $G \setminus G_{k-1}$.




Refined Canonical Order Example




Refined Canonical Order Example




Refined Canonical Order Example



Refined Canonical Order Example



Refined Canonical Order Example



Refined Canonical Order \rightarrow REL

We construct a REL as follows:

- For $i < j$, orient (v_i, v_j) from v_i to v_j ;
- v_k has incoming edges from v_{t_1}, \dots, v_{t_l} , we say that v_{t_1} is **left point** of v_k and v_{t_l} is **right point** of v_k .
- **Base edge** of v_k is (v_{t_a}, v_k) , where $t_a < k$ is minimal.
- If v_{k_1}, \dots, v_{k_o} are higher numbered neighbors of v_k , we call (v_k, v_{k_1}) **left edge** and (v_k, v_{k_o}) **right edge**.


Lemma 1.

A left edge or right edge cannot be a base edge.

Proof. Suppose left edge (v_k, v_{k_1}) is base edge of v_{k_1} .

Since G triangulated, $(v_{t_1}, v_{k_1}) \in E(G)$.

Contradiction since $v_k > v_{t_1}$.




Refined Canonical Order \rightarrow REL

Lemma 2.

An edge is either a **left edge**, a **right edge** or a **base edge**.

Proof.

- Exclusive “or” follows from Lemma 1.
- Let (v_{t_a}, v_k) be **base edge** of v_k .
- v_{t_a} is **right point** of $v_{t_{a-1}}$; $v_{t_i < a}$ is right point of $v_{t_{i-1}}$:
 - v_{t_i} has at least two higher-numbered neighbors.
 - One of them is v_k ; the other one is either $v_{t_{i-1}}$ or $v_{t_{i+1}}$.
 - For $1 \leq i < a - 1$, it is $v_{t_{i-1}}$.
- Analogously, $v_{t_{i \geq a}}$ is **left point** of $v_{t_{i+1}}$
- Edges (v_{t_i}, v_k) , $1 \leq i < a - 1$, are **right edges**.
- Similarly, (v_{t_i}, v_k) , for $a + 1 \leq i \leq l$, are **left edges**.



Refined Canonical Order \rightarrow REL

Coloring.

- Color **right** (**left**) edges in **red** (**blue**).
- Color a **base edge** (v_{t_i}, v_k) **red** if $i = 1$ and **blue** if $i = l$ and otherwise arbitrarily.



Let T_r be the red edges and T_b the blue edges.

Lemma 3.


$\{T_r, T_b\}$ is a regular edge labeling.

Proof.

$$k_o \geq 2$$




- $k_1 < k_2 < \dots < k_d$ and $k_d > k_{d+1} > \dots > k_o$
 - $(v_k, v_{k_i}), 2 \leq i \leq d-1$ are **blue**
 - $(v_k, v_{k_i}), d+1 \leq i \leq o-1$ are **red**
 - (v_k, v_{k_d}) is either **red** or **blue**
- \Rightarrow circular order of outgoing edges at v_k correct



Visualization of Graphs


Lecture 8:

Conact Representations of Planar Graphs: Triangle Contacts and Rectangular Duals




Part V: Computing the Coordinates


Jonathan Klawitter




From REL to st-digraphs to Coordinates




From REL to st-digraphs to Coordinates




From REL to st-digraphs to Coordinates




SN network G_{ver}
 dual of G_{ver}




From REL to st-digraphs to Coordinates




From REL to st-digraphs to Coordinates




From REL to st-digraphs to Coordinates



From REL to st-digraphs to Coordinates



From REL to st-digraphs to Coordinates




Rectangular Dual Algorithm

For a PTP graph $G = (V, E)$:

- Find a REL $\{T_r, T_b\}$ of G ;
- Construct a SN network G_{ver} of G (consists of T_b plus outer edges)
- Construct the dual G_{ver}^* of G_{ver} and compute a topological ordering f_{ver} of G_{ver}^*
- For each vertex $v \in V$, let g and h be the face on the left and face on the right of v . Set $x_1(v) = f_{\text{ver}}(g)$ and $x_2(v) = f_{\text{ver}}(h)$.
- Define $x_1(v_N) = 1, x_1(v_S) = 2$ and $x_2(v_N) = \max f_{\text{ver}} - 1, x_2(v_S) = \max f_{\text{ver}}$
- Analogously compute y_1 and y_2 with G_{hor} .
- For each $v \in V$, assign a rectangle $R(v)$ bounded by x-coordinates $x_1(v), x_2(v)$ and y-coordinates $y_1(v), y_2(v)$.

Reading off Coordinates to get Rectangular Dual



$$x_1(v_N) = 1, x_2(v_N) = 15$$

$$x_1(v_S) = 2, x_2(v_S) = 16$$

$$x_1(v_W) = 0, x_2(v_W) = 1$$

$$x_1(v_E) = 15, x_2(v_E) = 16$$

$$x_1(a) = 1, x_2(a) = 3$$

$$x_1(b) = 3, x_2(b) = 5$$

$$x_1(c) = 5, x_2(c) = 14$$

$$x_1(d) = 14, x_2(d) = 15$$

$$x_1(e) = 13, x_2(e) = 15$$

...

$$y_1(v_W) = 0, y_2(v_W) = 9$$

$$y_1(v_E) = 1, y_2(v_E) = 10$$

$$y_1(v_N) = 9, y_2(v_N) = 10$$


$$y_1(v_S) = 0, y_2(v_S) = 1$$

$$y_1(a) = 1, y_2(a) = 2$$

$$y_1(b) = 1, y_2(b) = 2$$

...

Reading off Coordinates to get Rectangular Dual



$$x_1(v_N) = 1, \quad x_2(v_N) = 15$$

$$x_1(v_S) = 2, \quad x_2(v_S) = 16$$

$$x_1(v_W) = 0, \quad x_2(v_W) = 1$$

$$x_1(v_E) = 15, \quad x_2(v_E) = 16$$

$$x_1(a) = 1, \quad x_2(a) = 3$$

$$x_1(b) = 3, \quad x_2(b) = 5$$

$$x_1(c) = 5, \quad x_2(c) = 14$$

$$x_1(d) = 14, \quad x_2(d) = 15$$

$$x_1(e) = 13, \quad x_2(e) = 15$$

...

$$y_1(v_W) = 0, \quad y_2(v_W) = 9$$

$$y_1(v_E) = 1, \quad y_2(v_E) = 10$$

$$y_1(v_N) = 9, \quad y_2(v_N) = 10$$

$$y_1(v_S) = 0, \quad y_2(v_S) = 1$$


$$y_1(a) = 1, \quad y_2(a) = 2$$

$$y_1(b) = 1, \quad y_2(b) = 2$$

...


Correctness of Algorithm (Sketch)

- If edge (u, v) exists, then $x_2(u) = x_1(v)$



$$x_2(u) = f_{\text{ver}}(g) = x_1(v)$$

- and the vertical segments of their rectangles overlap



$$\begin{aligned} y_1(v) = f_{\text{hor}}(a) &\leq y_1(u) = f_{\text{hor}}(b) \\ &< y_2(v) = f_{\text{hor}}(c) \leq y_2(u) = f_{\text{hor}}(d) \end{aligned}$$

- If path from u to v in red at least two edges long, then $x_2(u) < x_1(v)$.
- No two boxes overlap.
- for details see He's paper [He '93]

Rectangular Dual Result

Theorem.

Every PTP graph G has a rectangular dual, which can be computed in linear time.



Proof.

- Compute a planar embedding of G .
- Compute a refined canonical ordering of G .
- Traverse the graph and color the edges.
- Construct G_{ver} and G_{hor} .
- Construct their duals G_{ver}^* and G_{hor}^* .
- Compute a topological ordering for vertices of G_{ver}^* and G_{hor}^* .
- Assigning coordinates to the rectangles representing vertices.

Discussion


- A layout is **area-universal** if any assignment of areas to rectangles can be realized by a combinatorially equivalent rectangular layout.
- A rectangular layout is **area-universal** if and only if it is **one-sided**.
[Eppstein et al. SIAM J. Comp. 2012]

one-sided



not one-sided

- Area-universal **rectilinear** representation: possible for all planar graphs
- [Alam et al. 2013]: 8 sides (matches the lower bound)



Literature

Construction of triangle contact representations based on

- [de Fraysseix, de Mendez, Rosenstiehl '94] On Triangle Contact Graphs

Construction of rectangular dual based on

- [He '93] On Finding the Rectangular Duals of Planar Triangulated Graphs
- [Kant, He '94] Two algorithms for finding rectangular duals of planar graphs

and originally from

- [Kozłmiński, Kinnen '85] Rectangular Duals of Planar Graphs