


# Visualization of Graphs



## Lecture 3:

### Straight-Line Drawings of Planar Graphs I: Canonical Orderings and the Shift Method





Samuel Wolf

Summer term 2025



# Planar Graphs



$G$  is **planar**:


it can be drawn in such a way that no two edges intersect each other.

**planar embedding**:

clockwise orientation of adjacent vertices around each vertex

A planar graph can have many planar embeddings.

A planar embedding can have many planar drawings!



**faces**: Connected region of the plane bounded by edges

**Euler's polyhedra formula.**

$$\begin{array}{ccccccc} \# \text{faces} & - & \# \text{edges} & + & \# \text{vertices} & = & \# \text{conn.comp.} + 1 \\ f & - & m & + & n & = & c + 1 \end{array}$$

**Proof.** By induction on  $m$ :

$$m = 0 \Rightarrow f = 1 \text{ and } c = n \quad \checkmark$$

Induction hypothesis in  $G'$ :  
 $f' - m' + n' = c' + 1$

$$m \geq 1 \Rightarrow \text{delete some edge } e \Rightarrow m' = m - 1$$

$$\text{graph with edge } e \Rightarrow c' = c + 1$$

$$\text{graph with edge } e \Rightarrow f' = f - 1 \quad \checkmark$$

# Properties of Planar Graphs

## Euler's polyhedra formula.

$$\begin{array}{ccccccc} \# \text{faces} & - & \# \text{edges} & + & \# \text{vertices} & = & \# \text{conn.comp.} + 1 \\ f & - & m & + & n & = & c + 1 \end{array}$$

**Theorem.**  $G$  simple planar graph with  $n \geq 3$  vtc.

1.  $m \leq 3n - 6$
2.  $f \leq 2n - 4$
3. There is a vertex of degree at most 5.

**Proof.** 1. Every **edge** incident to  $\leq 2$  faces  
Every **face** incident to  $\geq 3$  edges

idea: count  
edge-face  
incidences

$$\Rightarrow 3f \leq \# \text{incidences} \leq 2m$$


$$\Rightarrow 6 \leq 3c + 3 = 3f - 3m + 3n \leq 2m - 3m + 3n = 3n - m$$

$$\Rightarrow m \leq 3n - 6$$

$$2. \quad 3f \leq 2m \leq 6n - 12 \Rightarrow f \leq 2n - 4$$

$$3. \quad \sum_{v \in V(G)} \deg(v) = 2m \leq 6n - 12$$

$$\Rightarrow \min_{v \in V(G)} \deg(v) \leq \text{average degree}(G) = \frac{1}{n} \sum_{v \in V(G)} \deg(v) \leq \frac{6n-12}{n} < 6$$



## Handshaking lemma.

$$\sum_{v \in V(G)} \deg(v) = 2|E(G)|.$$

# Triangulations

planar graph given with a planar embedding

A **plane (inner) triangulation** is a plane graph where every (inner) face is a triangle.


A **maximal planar graph** is a planar graph where adding any edge would violate planarity.

## Observation.

Any maximal plane graph is a plane triangulation (and vice versa).

## Lemma.


Any plane triangulation is 3-connected and thus has a unique planar embedding (up to mirroring).



We focus on plane triangulations:

## Lemma.

Every plane graph is a subgraph of a plane triangulation.



# Motivation

- Why planar and straight-line?

[Bennett, Ryall, Spaltzholz and Gooch '07]

## The Aesthetics of Graph Visualization

### 3.2. Edge Placement Heuristics

By far the most agreed-upon edge placement heuristic is to *minimize the number of edge crossings* in a graph [BMRW98, Har98, DH96, Pur02, TR05, TBB88]. The importance of avoiding edge crossings has also been extensively validated in terms of user preference and performance (see Section 4). Similarly, based on perceptual principles, it is beneficial to *minimize the number of edge bends* within a graph [Pur02, TR05, TBB88]. Edge bends make edges more difficult to follow because an edge with a sharp bend is more likely to be perceived as two separate objects. This leads to the heuristic of *keeping edge bends uniform* with respect to the bend's position on the edge and its angle [TR05]. If an edge must be bent to satisfy other aesthetic criteria, the angle of the bend should be as little as possible, and the bend placement should evenly divide the edge.

### Drawing conventions

- No crossings  $\Rightarrow$  planar
- No bends  $\Rightarrow$  straight-line


### Drawing aesthetics to optimize

- Area


# Towards Straight-Line Drawings

**Theorem.** [Kuratowski 1930]  
 $G$  planar  $\Leftrightarrow$   
 neither  $K_5$  nor  $K_{3,3}$  minor of  $G$

Kazimierz Kuratowski (1896–1980)



$K_5$



$K_{3,3}$

## Characterization

**Theorem.** [Hopcroft & Tarjan 1974]  
 Let  $G$  be a graph with  $n$  vertices. There is an  $\mathcal{O}(n)$ -time algorithm to test whether  $G$  is planar.

Also computes a planar embedding in  $\mathcal{O}(n)$  time.



John Edward Hopcroft (1939–)

[en.wikipedia.org/wiki/User:Shakespeare](https://en.wikipedia.org/wiki/User:Shakespeare)



Robert Endre Tarjan (1948–)

Renatokeshet, GFDL via Wikimedia

## Recognition

**Theorem.** [Wagner 1936, Fáry 1948, Stein 1951]  
 Every planar graph has a planar drawing  
 where the edges are straight-line segments.



Klaus Wagner (1910–2000)

Autor: Konrad Jacobs, wikipedia

## Drawing

The algorithms implied by these theorems produce drawings whose area is **not** bounded by any polynomial in  $n$ .




# Planar Straight-Line Drawings

**Theorem.** [De Fraysseix, Pach, Pollack '90]

Every  $n$ -vertex planar graph has a planar straight-line drawing of size  $(2n - 4) \times (n - 2)$ .

## Idea.

- Find a *canonical order*  $(v_1, \dots, v_n)$  of the vertices of a triangulation.
- Start with the single edge  $(v_1, v_2)$ . Let this be the graph  $G_2$ .
- Let  $k \in \{3, \dots, n\}$ . To obtain  $G_{k+1}$ , add  $v_{k+1}$  to  $G_k$  so that the neighbors of  $v_{k+1}$  are on the outer face of  $G_k$ .
- The neighbors of  $v_{k+1}$  in  $G_k$  form a path of length at least two.



**Theorem.** [Schnyder '90]

Every  $n$ -vertex planar graph has a planar straight-line drawing of size  $(n - 2) \times (n - 2)$ .

(next lecture)


# Canonical Order – Definition

## Definition.

Let  $G$  be a plane triangulation on  $n \geq 3$  vertices.


An ordering  $\pi = (v_1, v_2, \dots, v_n)$  of  $V(G)$  is a **canonical order** if the following conditions hold for each  $k \in \{3, 4, \dots, n\}$ :

- (C1) Vertices  $\{v_1, \dots, v_k\}$  induce a biconnected inner triangulation; call it  $G_k$ .
- (C2) Edge  $(v_1, v_2)$  belongs to the outer face of  $G_k$ .
- (C3) If  $k < n$  then vertex  $v_{k+1}$  lies in the outer face of  $G_k$ , and the neighbors of  $v_{k+1}$  form a path on the boundary of  $G_k$ .






# Canonical Order – Example



- (C1) Vertices  $\{v_1, \dots, v_k\}$  induce a biconnected inner triangulation; call it  $G_k$ .
- (C2) Edge  $(v_1, v_2)$  belongs to the outer face of  $G_k$ .
- (C3) If  $k < n$  then vertex  $v_{k+1}$  lies in the outer face of  $G_k$ , and the neighbors of  $v_{k+1}$  form a path on the boundary of  $G_k$ .


# Canonical Order – Example

- (C1) Vertices  $\{v_1, \dots, v_k\}$  induce a biconnected inner triangulation; call it  $G_k$ .
- (C2) Edge  $(v_1, v_2)$  belongs to the outer face of  $G_k$ .
- (C3) If  $k < n$  then vertex  $v_{k+1}$  lies in the outer face of  $G_k$ , and the neighbors of  $v_{k+1}$  form a path on the boundary of  $G_k$ .




# Canonical Order – Example

- (C1) Vertices  $\{v_1, \dots, v_k\}$  induce a biconnected inner triangulation; call it  $G_k$ .
- (C2) Edge  $(v_1, v_2)$  belongs to the outer face of  $G_k$ .
- (C3) If  $k < n$  then vertex  $v_{k+1}$  lies in the outer face of  $G_k$ , and the neighbors of  $v_{k+1}$  form a path on the boundary of  $G_k$ .




# Canonical Order – Example

- (C1) Vertices  $\{v_1, \dots, v_k\}$  induce a biconnected inner triangulation; call it  $G_k$ .
- (C2) Edge  $(v_1, v_2)$  belongs to the outer face of  $G_k$ .
- (C3) If  $k < n$  then vertex  $v_{k+1}$  lies in the outer face of  $G_k$ , and the neighbors of  $v_{k+1}$  form a path on the boundary of  $G_k$ .




# Canonical Order – Example



- (C1) Vertices  $\{v_1, \dots, v_k\}$  induce a biconnected inner triangulation; call it  $G_k$ .
- (C2) Edge  $(v_1, v_2)$  belongs to the outer face of  $G_k$ .
- (C3) If  $k < n$  then vertex  $v_{k+1}$  lies in the outer face of  $G_k$ , and the neighbors of  $v_{k+1}$  form a path on the boundary of  $G_k$ .


# Canonical Order – Example

- (C1) Vertices  $\{v_1, \dots, v_k\}$  induce a biconnected inner triangulation; call it  $G_k$ .
- (C2) Edge  $(v_1, v_2)$  belongs to the outer face of  $G_k$ .
- (C3) If  $k < n$  then vertex  $v_{k+1}$  lies in the outer face of  $G_k$ , and the neighbors of  $v_{k+1}$  form a path on the boundary of  $G_k$ .



*chord:*  
edge joining two  
non-adjacent  
vertices in a cycle


# Canonical Order – Example



- (C1) Vertices  $\{v_1, \dots, v_k\}$  induce a biconnected inner triangulation; call it  $G_k$ .
- (C2) Edge  $(v_1, v_2)$  belongs to the outer face of  $G_k$ .
- (C3) If  $k < n$  then vertex  $v_{k+1}$  lies in the outer face of  $G_k$ , and the neighbors of  $v_{k+1}$  form a path on the boundary of  $G_k$ .




# Canonical Order – Example



- (C1) Vertices  $\{v_1, \dots, v_k\}$  induce a biconnected inner triangulation; call it  $G_k$ .
- (C2) Edge  $(v_1, v_2)$  belongs to the outer face of  $G_k$ .
- (C3) If  $k < n$  then vertex  $v_{k+1}$  lies in the outer face of  $G_k$ , and the neighbors of  $v_{k+1}$  form a path on the boundary of  $G_k$ .


# Canonical Order – Example

- (C1) Vertices  $\{v_1, \dots, v_k\}$  induce a biconnected inner triangulation; call it  $G_k$ .
- (C2) Edge  $(v_1, v_2)$  belongs to the outer face of  $G_k$ .
- (C3) If  $k < n$  then vertex  $v_{k+1}$  lies in the outer face of  $G_k$ , and the neighbors of  $v_{k+1}$  form a path on the boundary of  $G_k$ .




# Canonical Order – Example

- (C1) Vertices  $\{v_1, \dots, v_k\}$  induce a biconnected inner triangulation; call it  $G_k$ .
- (C2) Edge  $(v_1, v_2)$  belongs to the outer face of  $G_k$ .
- (C3) If  $k < n$  then vertex  $v_{k+1}$  lies in the outer face of  $G_k$ , and the neighbors of  $v_{k+1}$  form a path on the boundary of  $G_k$ .



# Canonical Order – Example



- (C1) Vertices  $\{v_1, \dots, v_k\}$  induce a biconnected inner triangulation; call it  $G_k$ .
- (C2) Edge  $(v_1, v_2)$  belongs to the outer face of  $G_k$ .
- (C3) If  $k < n$  then vertex  $v_{k+1}$  lies in the outer face of  $G_k$ , and the neighbors of  $v_{k+1}$  form a path on the boundary of  $G_k$ .

# Canonical Order – Existence

## Lemma.



Every plane triangulation has a canonical order.

Consider any  $n$ -vertex plane triangulation. We show this statement by induction on  $k$  from  $n$  down to 3.

**Induction base ( $k = n$ ):** Let  $G_n = G$ , and let  $v_1, v_2, v_n$  be the vertices of the outer face of  $G_n$ . Conditions (C1)–(C3) hold.

**Induction hypothesis:** Vertices  $v_{n-1}, \dots, v_{k+1}$  have been chosen such that conditions (C1)–(C3) hold for every  $i \in \{k+1, \dots, n\}$ .

**Induction step:** Consider  $G_k$ . We search for  $v_k$ .



- (C1)  $G_k$  biconnected inner triangulation
- (C2)  $(v_1, v_2)$  on outer face of  $G_k$
- (C3)  $k < n \Rightarrow v_{k+1}$  in outer face of  $G_k$ , neighbors of  $v_{k+1}$  form path on boundary of  $G_k$


We need to show:

1.  $v_k$  not incident to **chord** is sufficient.
2. Such  $v_k$  exists.

# Canonical Order – Existence


## Claim 1.

If  $v_k$  is not incident to a **chord**, then  $G_{k-1}$  is biconnected.



## Claim 2.

There exists a vertex in  $G_k$  that is not incident to a chord as choice for  $v_k$ .



This completes the proof of the lemma.  $\square$

# Canonical Order – Implementation

outer face

CanonicalOrder( $G, \langle v_1, v_2, v_n \rangle$ )

**foreach**  $v \in V(G)$  **do**

└  $\text{chords}(v) \leftarrow 0$ ;  $\text{out}(v) \leftarrow \text{false}$ ;  $\text{mark}(v) \leftarrow \text{false}$

$\text{out}(v_1), \text{out}(v_2), \text{out}(v_n) \leftarrow \text{true}$

**for**  $k = n$  **downto** 3 **do**

choose  $v \in V(G) \setminus \{v_1, v_2\}$  such that  $\text{mark}(v) = \text{false}$ ,  
 $\text{out}(v) = \text{true}$ ,  $\text{chords}(v) = 0$  // use list of candidates

$v_k \leftarrow v$ ;  $\text{mark}(v_k) \leftarrow \text{true}$ ;  $\text{out}(v_k) \leftarrow \text{false}$


let  $w_p, \dots, w_q$  be the ordered unmarked neighbors of  $v_k$

**for**  $i = p + 1$  **to**  $q - 1$  **do** //  $O(n)$  time in total  
 └  $\text{out}(w_i) \leftarrow \text{true}$  //  $O(m) = O(n)$  in total

**foreach**  $u \in \text{Adj}[w_i] \setminus \{w_{i-1}, w_{i+1}\}$  **do**  $\leftarrow$   
 └ **if**  $\text{out}(u)$  **then**  $\text{chords}(w_i)++$ ,  $\text{chords}(u)++$

**if**  $p + 1 = q$  **then**  $\text{chords}(w_p)--$ ,  $\text{chords}(w_q)--$

- $\text{chord}(v) =$   
 # chords incident to  $v$
- $\text{out}(v) = \text{true}$  iff  $v$  on boundary of current outer face
- $\text{mark}(v) = \text{true}$  iff  $v$  has received a number  $\geq k$



## Lemma.

Algorithm CanonicalOrder computes a canonical order of a plane graph in  $\mathcal{O}(n)$  time.




# Shift Method – Idea

## Drawing invariants:

$G_k$  is drawn such that

- $v_1$  is at  $(0, 0)$ ,  $v_2$  is at  $(2k - 4, 0)$ ,
- boundary of  $G_k$  (minus edge  $\{v_1, v_2\}$ ) is drawn x-monotone,
- each edge on the boundary of  $G_k$  (except  $\{v_1, v_2\}$ ) is drawn with slopes  $\pm 1$ .




# Shift Method – Idea

## Drawing invariants:

$G_k$  is drawn such that

- $v_1$  is at  $(0, 0)$ ,  $v_2$  is at  $(2k - 4, 0)$ ,
- boundary of  $G_k$  (minus edge  $\{v_1, v_2\}$ ) is drawn x-monotone,
- each edge on the boundary of  $G_k$  (except  $\{v_1, v_2\}$ ) is drawn with slopes  $\pm 1$ .




# Shift Method – Idea

## Drawing invariants:


$G_k$  is drawn such that

- $v_1$  is at  $(0, 0)$ ,  $v_2$  is at  $(2k - 4, 0)$ ,
- boundary of  $G_k$  (minus edge  $\{v_1, v_2\}$ ) is drawn x-monotone,
- each edge on the boundary of  $G_k$  (except  $\{v_1, v_2\}$ ) is drawn with slopes  $\pm 1$ .




Will  $v_{k+1}$  lie on the grid?






Yes, because  $w_p$  and  $w_q$  have even Manhattan distance  $\Delta x + \Delta y$ .






# Shift Method – Example






# Shift Method – Example



# Shift Method – Example






# Shift Method – Example











# Shift Method – Example






# Shift Method – Example






# Shift Method – Example






# Shift Method – Example






# Shift Method – Example



# Shift Method – Example






# Shift Method – Example





# Shift Method – Example




# Shift Method – Planarity

## Observations.

- Each internal vertex is **covered** exactly once.
- Covering relation defines a tree in  $G$
- and a forest in  $G_i$ ,  $1 \leq i \leq n - 1$ .

## Lemma.

Let  $0 \leq \delta_1 \leq \delta_2 \leq \dots \leq \delta_t \in \mathbb{N}$ , s.t.  $\delta_{p+1} - \delta_p \geq 1$ ,  $\delta_q - \delta_{q-1} \geq 1$ ,  $\delta_q - \delta_p \geq 2$  and even. If we shift  $L(w_i)$  by  $\delta_i$  to the right, then we get a planar straight-line drawing.




## Proof by induction:

If  $G_{k-1}$  is drawn planar and straight-line, then so is  $G_k$ .

Ideas:

- New edges don't intersect other edges ( $\rightarrow$  invariants).
- Edges within each  $L(w_i)$  do not change.
- Other edges lie within triangles that only become flatter without causing new intersections.




# Shift Method – Pseudocode

canonical order of  $V(G)$

ShiftMethod( $G, (v_1, v_2, \dots, v_n)$ )

**for**  $k = 1$  to 3 **do**

$L(v_k) \leftarrow \{v_k\}$

$P(v_1) \leftarrow (0, 0); P(v_2) \leftarrow (2, 0), P(v_3) \leftarrow (1, 1)$  

**for**  $k = 4$  to  $n$  **do**

    Let  $\partial G_{k-1}$  be  $v_1 = w_1, w_2, \dots, w_{t-1}, w_t = v_2$ .

    Let  $w_p, \dots, w_q$  be the neighbors of  $v_k$ .

**foreach**  $v \in \bigcup_{i=p+1}^{q-1} L(w_i)$  **do**      *//  $\mathcal{O}(n^2)$  in total*

$x(v) \leftarrow x(v) + 1$


**foreach**  $v \in \bigcup_{i=q}^t L(w_i)$  **do**      *//  $\mathcal{O}(n^2)$  in total*

$x(v) \leftarrow x(v) + 2$

$P(v_k) \leftarrow$  intersection of slope- $\pm 1$  diagonals  
        through  $P(w_p)$  and  $P(w_q)$

$L(v_k) \leftarrow \bigcup_{i=p+1}^{q-1} L(w_i) \cup \{v_k\}$

**return**  $P(v_1), \dots, P(v_n)$



**Running Time?**

# Shift Method – Linear-Time Implementation


## Idea 1.

To compute  $x(v_k)$  and  $y(v_k)$ ,  
we need only  $y(w_p)$ ,  $y(w_q)$ , and  $x(w_q) - x(w_p)$

## Idea 2.

Instead of storing explicit x-coordinates,  
we store, for each vertex within a specific spanning tree,  
the x-distance to its parent ( $v_1$  is the root).

After an x-distance is computed for each  $v_k$ ,  
use preorder traversal to compute all x-coordinates.



- (1)  $x(v_k) = \frac{1}{2}(x(w_q) + x(w_p) + y(w_q) - y(w_p))$
- (2)  $y(v_k) = \frac{1}{2}(x(w_q) - x(w_p) + y(w_q) + y(w_p))$
- (3)  $x(v_k) - x(w_p) = \frac{1}{2}(x(w_q) - x(w_p) + y(w_q) - y(w_p))$

# Shift Method – Linear-Time Implementation


## Relative x-distance tree.

For each vertex  $v$  store

- x-offset  $\Delta_x(v)$  from parent
- y-coordinate  $y(v)$

## Calculations.

- $\Delta_x(w_{p+1})++$ ,  $\Delta_x(w_q)++$
- $\Delta_x(w_p, w_q) = \Delta_x(w_{p+1}) + \dots + \Delta_x(w_q)$
- $\Delta_x(v_k)$  by (3)      ■  $y(v_k)$  by (2)
- $\Delta_x(w_q) = \Delta_x(w_p, w_q) - \Delta_x(v_k)$
- $\Delta_x(w_{p+1}) = \Delta_x(w_{p+1}) - \Delta_x(v_k)$



takes  $\mathcal{O}(n)$  time in total 😊

$$(1) \quad x(v_k) = \frac{1}{2}(x(w_q) + x(w_p) + y(w_q) - y(w_p))$$

$$(2) \quad y(v_k) = \frac{1}{2}(x(w_q) - x(w_p) + y(w_q) + y(w_p))$$

$$(3) \quad \underbrace{x(v_k) - x(w_p)}_{\Delta_x(v_k)} = \frac{1}{2}(\underbrace{x(w_q) - x(w_p)}_{\Delta_x(w_p, w_q)} + y(w_q) - y(w_p))$$

# Discussion

- The shift method by de Fraysseix, Pach, and Pollack provides an algorithmic tool to efficiently draw a plane graph onto a polynomial-size grid using only straight-line edges.
- The linear-time implementation was later proposed by Chrobak and Payne.
- Although we are guaranteed to get a very small grid, only straight-line edges, and no edge crossings, the resulting drawings are not always visually pleasing: the drawings tend to have very small angles and a big variance in the size of the triangular faces.
- A quite different approach yielding similar results is by Schnyder (→ next lecture).

# Literature

- [PGD Ch. 4.2] for detailed explanation of the shift method
- [de Fraysseix, Pach, Pollack 1990] “How to draw a planar graph on a grid”
  - original paper introducing the shift method
- [Chrobak, Payne 1995] “A linear-time algorithm for drawing a planar graph on a grid”
  - original paper on how to implement the shift method in linear time